


**4-6 Chemistry /5-6 Trilogy – Rate and extent of chemical change**

- 1.0** A student heated hydrated cobalt chloride.  
The word equation shows the reaction.



- 1.1** The student recorded some observations from this experiment.  
Suggest **two** observations the student may have written down.

[2 marks]

---



---

- 1.2** The student added anhydrous cobalt chloride to water and measured the temperature rise.  
The student's results are shown in the table below.

|                               | <b>Trial 1</b> | <b>Trial 2</b> | <b>Trial 3</b> |
|-------------------------------|----------------|----------------|----------------|
| <b>Temperature rise in °C</b> | 9.5            | 9.2            | 9.2            |

Calculate the mean temperature rise.

[1 mark]

---


Temperature = \_\_\_\_\_ °C

- 1.3** During the reaction in **1.2**, the temperature increased.  
Name the type of reaction that causes the temperature to rise.

[1 mark]

---

**2.0** A student investigated the effect of temperature on the rate of a reaction.  
**Figure 1** below shows the apparatus the student used.



**2.1** Name a piece of apparatus which could be used to measure the volume of the acid. [1 mark]

\_\_\_\_\_

**2.2** The reaction forms a precipitate.  
 When should the student stop timing the reaction? [1 mark]

\_\_\_\_\_

**2.3** State the dependent and independent variables in the investigation. [2 marks]

Dependent \_\_\_\_\_

Independent \_\_\_\_\_

**2.4** The student only carried out each test once.  
 Explain why repeating the experiment would improve the results. [1 mark]


\_\_\_\_\_

**2.5** Describe how a preliminary investigation could be used to find an appropriate temperature range. [2 marks]

\_\_\_\_\_

\_\_\_\_\_

- 2.6 Another student used a different experiment to investigate the rate of reaction. This student measured the loss of mass every minute. The student's results are shown in **Graph 1** below:



Add labels to the graph to show:

- when the reaction is complete
- when the rate of reaction is fastest
- when half the reactants have been used up.

[3 marks]

3.0 A student investigated how the concentration of hydrochloric acid affected the rate of reaction between hydrochloric acid (HCl) and magnesium ribbon to produce magnesium chloride (MgCl<sub>2</sub>) and hydrogen (H<sub>2</sub>).

3.1 Complete and balance the equation for the reaction:

[2 marks]




Figure 2 below shows the apparatus the student used.




Table 1 shows the results of the experiment.

Table 1

| Concentration of hydrochloric acid in mol/dm <sup>3</sup> | Time taken for 30 cm <sup>3</sup> of hydrogen to be produced in s |         |         |      | Mean rate of reaction in cm <sup>3</sup> /s |
|---|---|---------|---------|------|---|
|   | Trial 1   | Trial 2 | Trial 3 | Mean |   |
| 0.4   | 158   | 150     | 154     | 154  | 0.19  |
| 0.8   | 77  | 77      | 74      | 76   | 0.39  |
| 1.2   | 68  | 51      | 49      |      |   |
| 1.6   | 37  | 39      | 38      | 38   | 0.79  |
| 2.0   | 30  | 29      | 31      | 30   | 1.00  |

**3.2** Calculate the rate of reaction when 1.2 mol/dm<sup>3</sup> hydrochloric acid is added to magnesium.

Use the equation below.

$$\text{mean rate of reaction} = \frac{\text{volume of gas in cm}^3}{\text{mean time taken in s}}$$

[3 marks]

Mean rate of reaction = \_\_\_\_\_ cm<sup>3</sup>/s

**3.3** Give **two** variables which the student should control during this investigation.

[2 marks]

---




---

**3.4** On **Figure 3**, use the results from **Table 1** to

- plot a graph of rate of reaction and concentration of acid
- draw a best fit line

[3 marks]

**Figure 3**



**3.5** Using the idea of particle collisions, explain why the reaction rate is faster when the concentration of the acid is greater.

[2 marks]

---

---

**3.6** The student used magnesium ribbon.  
State a change that could be made to the magnesium to speed up the reaction.

[1 mark]

---

**3.7** Explain in terms of the particles why the change you gave in **3.6** would increase the speed of reaction.

[1 mark]

---

4.0 This question is about reversible reactions and chemical equilibrium.

4.1 Reversible reactions can reach equilibrium in a closed system.  
What is meant by a **closed system**?

[1 mark]

---

4.2 Explain why a reaction seems to have finished when a reversible reaction reaches equilibrium.

[2 marks]

---




---

Ethanol can be produced in a reversible reaction from ethene and steam.  
The equation for the reaction is:



**Figure 4** shows the reaction profile for the reaction.

**Figure 4**



4.3 How does the diagram show that the reaction is exothermic?

[1 mark]

---

**4.4** A catalyst can be used for the reaction.

Indicate on **Figure 4**:

- the reaction profile for a catalysed reaction
- the activation energy for a catalysed reaction.

[2 marks]

**4.5** State what is meant by **activation energy**.

[1 mark]

---

---

**4.6** Give one similarity and one difference in the energy transfer for the back reaction to form ethene and water from ethanol.

[2 marks]


Similarity: \_\_\_\_\_

Difference: \_\_\_\_\_



4.7 A company manufactures ethanol (C<sub>2</sub>H<sub>5</sub>OH).

The reaction for the process is:



The temperature and pressure can be changed to increase the yield of ethanol at equilibrium.

The forward reaction is exothermic.

The conditions used in the process are:

- 60 atmospheres pressure
- 200 °C
- phosphoric acid catalyst.

Explain why these conditions are used in this process.

Use the equation and your knowledge of reversible reactions

Consider **both** yield **and** rate of reaction in your answer.

[6 marks]

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---

---

---

MARK SCHEME

| Qu No. |                                   | Extra Information | Marks |
|--------|-----------------------------------|-------------------|-------|
| 1.1    | (solid) changes from pink to blue |                   | 1     |
|        | Droplets of water / steam         |                   | 1     |
| 1.2    | 9.3 °C                            |                   | 1     |
| 1.3    | Exothermic                        |                   | 1     |

| Qu No. |  | Extra Information  | Marks                      |
|--------|--|--|----------------------------|
| 2.1    | Measuring cylinder   | Allow burette/pipette  | 1                          |
| 2.2    | When the cross cannot be seen through the solution   | ignore when the solution is cloudy   | 1                          |
| 2.3    | (dependent)<br>Time taken for the cross to disappear   |  | 1                          |
|        | (independent)<br>Temperature   |  | 1                          |
| 2.4    | To check the results.<br>So you know the readings are accurate.<br>To eliminate/ignore anomalous results.  | Allow to improve reliability.  | 1                          |
| 2.5    | Two temperatures are suggested that constitute a range   |  | 1                          |
|        | Understanding demonstrated that an appropriate range will allow a pattern or trend to be seen in the results   |  | 1                          |
| 2.6    | <p style="text-align: center;"><b>Graph 1</b></p>  <p>A: reaction is complete<br/>B: reaction is fastest<br/>C: half the reactants have been used up.</p> | <p>A: Must be after graph levels off</p> <p>B: Any point on straight line up before it changes gradient</p> <p>C: When loss of mass is 20g</p> | <p>1</p> <p>1</p> <p>1</p> |

| Qu No. |  | Extra Information   | Marks |
|--------|--|---|-------|
| 3.1    | Formulae in correct place  |   | 1     |
|        | Correct balancing  | Allow 2 marks for<br>$\text{Mg(s)} + 2\text{HCl(aq)} \rightarrow \text{MgCl}_2(\text{aq}) + \text{H}_2(\text{g})$         | 1     |
| 3.2    | $(49+51)/2$  |   | 1     |
|        | (mean =) 50  | Allow 2 marks for 50 without working  | 1     |
|        | $(30/50 =) 0.60$   | Allow 2 marks for 0.54 where anomaly has been included in mean  | 1     |
| 3.3    | any <b>two</b> from: <ul style="list-style-type: none"> <li>• volume of acid</li> <li>• temperature (of acid)</li> <li>• length of magnesium (ribbon)</li> </ul> | Do not allow concentration of acid<br><br>allow mass of magnesium ribbon  | 2     |
| 3.4    | All points plotted correctly   | $\pm \frac{1}{2}$ small square<br>Allow 1 mark for 4 plotted correctly<br><br>Allow ecf for anomalous point at (1.2,0.54) | 2     |
|        | Best fit straight line   | Should not be influenced by anomaly   | 1     |
| 3.5    | Particles must collide in order to react   |   | 1     |
|        | Collision frequency increases as concentration increases   |   | 1     |
| 3.6    | cut it up <b>or</b> increase the surface area  | Allow grind it up <b>or</b> make a powder<br><br>do <b>not</b> accept make it smaller <b>or</b> use a smaller piece       | 1     |
| 3.7    | Reference to particle theory eg more collisions between acid ions/particles and atoms/particles of magnesium   |   | 1     |

| Qu No. |  | Extra Information   | Marks      |
|--------|--|---|------------|
| 4.1    | nothing can enter and nothing can leave the reaction   | allow sealed reaction vessel  | 1          |
| 4.2    | at equilibrium the forward and backward reactions have same rate<br><br>so there is no (overall) change in quantities of reactants and products                            |   | 1<br><br>1 |
| 4.3    | the products are at a lower energy level than the reactants  | accept products have less energy or less energy at the end than the beginning                   | 1          |
| 4.4    | Pathway drawn from reactants to products, below original pathway<br><br>Indication of activation energy from reactant level to highest point on catalysed reaction pathway |   | 1<br><br>1 |
| 4.5    | Minimum amount of energy needed by particles to react  |   | 1          |
| 4.6    | <i>Similarity</i><br>Same amount of energy transferred<br><br><i>Difference</i><br>Endothermic reaction  | Allow 45 kJ of energy transferred (given in 4.7 below)<br><br>Allow energy taken in by reaction | 1<br><br>1 |

|   |   |     |
|---|---|-----|
| 4.7   |   |     |
| <b>Level 3:</b>   | A detailed and coherent explanation is given, which demonstrates a broad understanding of the key scientific ideas. The response makes logical links between the points raised and uses sufficient examples to support these links. | 5-6 |
| <b>Level 2:</b>   | An explanation is given which demonstrates a reasonable understanding of the key scientific ideas. Links are made but may not be fully articulated and / or precise.  | 3-4 |
| <b>Level 1:</b>   | Simple statements are made which demonstrate a basic understanding of some of the relevant ideas. The response may fail to make logical links between the points raised.  | 1-2 |
|   | No relevant content   | 0   |
| <b>Indicative content</b>   |   |     |
| <p><b>60 atmospheres pressure</b></p> <ul style="list-style-type: none"> <li>• high pressure gives a high yield of ethanol</li> <li>• too high a pressure causes risk of explosion</li> <li>• high pressure costly to maintain</li> <li>• a high pressure will cause the rate to be higher</li> <li>• 2 moles of gas become 1 (or fewer moles of gas in products)</li> </ul> <p><b>200 °C</b></p> <ul style="list-style-type: none"> <li>• high temperature increases the rate of reaction</li> <li>• optimum temperature</li> <li>• (forward reaction is exothermic so) a high yield of ethanol requires a low temperature</li> <li>• but too low a temperature causes the rate of reaction to be too slow</li> </ul> <p><b>phosphoric acid catalyst</b></p> <ul style="list-style-type: none"> <li>• a catalyst speeds up the reaction<br/>a phosphoric acid catalyst allows a lower temperature to be used (saving energy and causing a higher yield)</li> <li>• phosphoric acid catalyst increases the rate of reaction equally in both reactions</li> </ul> <p><b>others</b></p> <ul style="list-style-type: none"> <li>• compromise conditions</li> <li>• unreacted ethene and steam is recycled</li> </ul> |   |     |